

AMERICAN JOURNAL OF PHARMACY AND THE SCIENCES SUPPORTING PUBLIC HEALTH

Since 1825

COMMITTEE ON PUBLICATION

E. Fullerton Cook, Sc. D., Ph. M. Mitchell Bernstein, M. D. J. W. Sturmer, D. Sc.
John K. Thum, Ph. M. Louis Gershenfeld, P. D., Ph. M. Joseph W. E. Harrison, Ph. M.

IVOR GRIFFITH, Ph. M., Sc. D., Editor
Linwood F. Tice, M. Sc., Assistant to the Editor
John E. Kramer, B. Sc., Business Manager

Vol. 111.

DECEMBER, 1939

No. 12

CONTENTS

Editorial:

The Position of Pharmacy in Public Health 462

Original Article:

A New Approach to the Standardization of Digitalis. By A. Viehoveer,
N. H. Sokoloff and A. A. Taransky 466

Abstracts from, and Reviews of, the Literature of the Sciences Sup-
porting Public Health 475

Solid Extracts 480

Book Review 484

Index to Volume 111 485

Annual Subscription, \$3.00

Foreign Postage, 25 Cents Extra

Single Numbers, 30 Cents.

Back Numbers, 50 Cents

Entered as Second-Class Matter at the Post Office at Philadelphia, Pa.,
Under the Act of March 3, 1879

Acceptance for Mailing at Special Rate of Postage Provided for in Section 1103
Act of October 3, 1917 Authorized February 15, 1920.

E D I T O R I A L

On these pages the editor offers his opinions, unshackled by advertising patrons and unrestrained by anything save a sense of the decent and the truthful. The editor, alone, is responsible for their type, their tone and their tenor.

THE POSITION OF PHARMACY IN PUBLIC HEALTH

PHARMACY'S history is one of its proudest assets, for "the art of the apothecary" has been practiced since the first man had his first illness. Thus, when our ancestors fell heir to, or chose their duties, some of them, by nature or by inclination, came into the practice of medicine and pharmacy.

Through many centuries, the two professions, were interdependent and inseparable. In most instances medicine and pharmacy were practiced simultaneously by the same individuals. Then, in the twelfth and thirteenth centuries, further delineation of communal duties brought about the establishment of apothecary shops, or drug stores, owned and managed by men whose specialized work as pharmacists demanded so much of their time that they could not enter also into the practice of medicine.

Today we find the same distinction of duties, together with harmony and understanding between the professions of medicine and pharmacy, each, in its own way and with the assistance of the other, playing its part in the never ending struggle against pain, sickness and disease. Dentistry, nursing and the other public health services are indispensable allies in this united endeavor to bring about longer and healthier lives. It has been a progressive battle. In a period of only a century and a half in this country, man's life expectancy has risen from 35 to 62 years. The products provided by modern pharmacy have contributed in no small measure to this result.

The Meaning of the Profession

Pharmacy is the art and science of preparing from natural and synthetic sources suitable and convenient materials for use as drugs. It also comprehends the compounding of drugs and the dispensing of drugs and medicines according to prescription, and their distribution in other ways. Pharmacy also embraces the collection, identification,

preservation, analysis and standardization of drugs and medicines; the synthesis of medicinal chemicals and the preparation of biological products. In short, the primary function of pharmacy is to prepare medicines for those who need them.

Pharmacy's Progress

From its domination in early days by alchemy and superstition, pharmacy has developed into a highly specialized art involving the application of the exact and exacting sciences. The necessity of trade has always been a more or less important factor in its existence, and has, at times, seemed to obscure its real value and function. Yet, through a happy fellowship of professionalism and commercialism, the ready availability and distribution of drugs and medicines throughout this country is provided for. In the United States there are a sufficient number of adequately trained registered pharmacists. One hundred and twenty thousand of these professional and scientific workers, practicing in 60,000 registered pharmacies, and 3000 hospitals, insure that no person is at any time very far removed from an adequate and reliable source of medicaments.

Based on the country's population, this indicates one drug store ready to serve each 2000 people. In the aggregate, 200,000,000 prescriptions are filled by these pharmacists each year. Hospital and clinic pharmacies fill countless others.

The Pharmacist Serves the Public Welfare

In addition to the compounding of prescriptions, retail pharmacists in drug stores render many other useful services. Through the knowledge which they have gained in years spent in colleges of pharmacy in their apprenticeships and in their practice, they make a special contribution to the welfare of the citizen. Their intelligence in the methods of preparing and properly combining medicinal substances places them constantly in a position to assist practicing physicians in the successful treatment of disease. They likewise supply physicians and dentists with sterile solutions, antiseptics, surgical aids, anesthetics, stains used in the study of bacteria; also various other materials used in diagnosis and treatment.

The retail pharmacist is also a frequent source of supply for veterinary medicines, insecticides, fungicides, rodent poisons, household disinfectants and many similar articles with which he has become so familiar.

Serums, and vaccines, foods for infants and invalids, sickroom and first-aid supplies, and sanitation aids are customarily distributed by pharmacists because of the specialized knowledge required for the preparation, storage and use of these items.

The retail or hospital pharmacist frequently manufactures such compounds as must be freshly prepared, and he also manufactures a number of medicines which can be economically prepared in his own laboratory. He is also the center of distribution and the focal point of information and advice concerning the myriad of vital preparations manufactured in the great, centrally located laboratories.

And here we have a modern phase of pharmacy, to the casual on-looker little known, because part of the practice of the profession has passed from the hands of the "apothecary" into other yet capable hands. The extensive drug industry—huge drug collecting organizations and milling concerns, large manufacturers of organic and inorganic chemicals of medical value, immense pharmaceutical manufacturing establishments, biologic laboratories, institutes of pharmaceutical research—all of these are integral parts of the pharmacy of today, with the corner Drug Store an important link in the chain of agencies necessary to provide all the preparations required by modern medicine. As the actual prescription-compounding, medicine-supplying station, the corner drug store, notwithstanding its side lines, is a valuable, serviceable spot in every community.

The Pharmacist in the Community

As a general rule, pharmacists are intensely civic-minded. They are often active in various service clubs, chambers of commerce, church and charity organizations, and other groups working for the betterment of the community. Often, their stores are the centers of neighborhood activity along these lines. Open for a greater part of the day than most business establishments, they lend themselves admirably to this purpose.

The Law Recognizes the Pharmacist

Recognized as a public health profession in every civilized country in the world, the practice of pharmacy in all such countries is regulated by law. In the United States the laws of the various States limit the practice of pharmacy to those who are properly qualified and licensed as determined by formal collegiate education and subsequent state governmental examination. These same laws have placed the

legal distribution of poisons, potent drugs and medicines and narcotics under the supervision of pharmacists.

Such laws are just, and work for public benefit. Under these regulations, pharmacy is able to render an incomparable service to the community. If the public interest be given full consideration there should be no curtailment of the services now rendered by the pharmacists of the nation. Obviously, therefore, any effort in the direction of concentrating pharmaceutical facilities in governmental departments or health centers except in a few very special cases, would reduce the availability of adequate assistance on a private scale. Pharmacy is a profession of service to each citizen in each country. It must and will always be so.

Your Pharmacist Greets You

Whether you are well, or whether you are ill, your pharmacist greets you. He is eminently prepared to render professional service without limit. He is happy and willing to render the multitude of other services you are accustomed to expect of your favorite pharmacy.

Your pharmacist is your friend.

Effect of Radiant Energy on Thermophilic Organisms in Sugar. H. H. Hall and J. C. Keane. *J. Ind. & Eng. Chem.* 31, 1168 (1939). Results obtained by the Bureau of Agricultural Chemistry indicate that the spores of a thermophilic organism which spoils canned food, *B. stearothermophilus* Donk, are killed in dry white sugar by radiant energy rays most of which are in the region 253.7μ . The lethal action is enhanced when the sugar crystals are kept in constant motion in the sugar granulator during irradiation. For this purpose, 24 thirty inch lamps were installed in the sugar outlet end of a granulator. Under these conditions an average of 47.8 per cent. of the spores were killed by irradiating eight successive batches of sugar. No chemical changes were noted in the irradiated product. The results indicate the possibility of sterilizing sugar and other such substances by this method.

L. F. T.

ORIGINAL ARTICLES

A NEW APPROACH TO THE STANDARDIZATION OF DIGITALIS

By Arno Viehoveer, N. H. Sokoloff and A. A. Taransky * †

I. Introduction

THE unsolved problem of devising a reliable method for the standardization of digitalis and its preparations has been intensively investigated for several decades. Although it is agreed that graded dosage of this important heart tonic can be permitted only by standardization, the majority of methods devised for measuring potency are in reality merely toxicity assays. Were it true that the toxicity of a digitalis preparation is related to its therapeutic efficiency in correcting cardiac dysfunction, then such methods for the assay of potency would have a logical basis. However, we believe that potency in the case of digitalis can scarcely be considered as synonymous with toxicity. Thus, the official methods of standardization are toxicity assays which indicate the margin of safety of a preparation and not the actual therapeutic value.

Significant discrepancies in results have been reported by different workers using the same methods for a single sample of digitalis, and with the use of other methods the disagreement has been more marked. Even in the case of the U. S. P. XI Digitalis Standard, Rowe has reported that there is a variable action, and that it is 50 per cent. stronger than the U. S. P. X Standard (8).

The senior author and his previous students (5 and 6) had studied the effect of digitalis and certain of its constituents upon the heart of *Daphnia magna*. Digitoxin rapidly depressed the rate of heartbeat. Also, as recorded in micromovies (7), the spastic heartbeat of daphnia produced by chloroform was abolished, and a regular rhythm with increased force established through the use of digitalis and Verodigen—a preparation of digitalis glucosides. More recent studies have disclosed the fact that certain types of drug-induced depression of the rate of heartbeat could be wholly or partially prevented

* From the Gross Laboratory for Biological and Biochemical Research.

† We are indebted to Dr. Isadore Cohen for invaluable aid and advice in these experiments.



in the presence of an active digitalis preparation. These findings suggested an approach to a new method of studying and perhaps standardizing digitalis by its desirable remedial activity.

II. Description and Culture of *Daphnia Magna*

Daphnia magna is a small (1/10 inch) transparent, fresh-water crustacean belonging to the Cladocera. It possesses well developed muscular, nervous and glandular systems. The ease of observation of the internal organs functioning normally and under drug or toxic action permits direct quantitative measurements (3). The propagation of *Daphnia magna* has been reported by Viehoveer (1) and the methods of standardization by Viehoveer and Cohen (4). Cultured under uniform conditions, *Daphnia magna* exhibits a definite rhythm and periodicity of growth. Reproduction is chiefly through diploid parthenogenesis. Numerous offspring of genetical uniformity is thus assured for statistical studies. The offspring can reproduce in ten days, while the life span of the species at 70 degrees F. is approximately ninety days.

III. Description of the Methods Used in the *Daphnia* Experiments

In the earlier experiments the hanging-drop technique was utilized. Double depression slides were found to be especially satisfactory since a control animal could be mounted for direct comparison with the test animal which was suspended in the drug mixture. These mixtures consisted either of yohimbine (2), veratrine or atropine which depress the rate of heartbeat of daphnia, and digitalis or digitalis glucosides in varying proportions. Although this method possessed the virtue of simplicity, there were numerous disadvantages and sources of errors. The amount of the test solution was limited to a single drop, and there was an unavoidable variation in the size of the drop from one experiment to another. The cooling effect of evaporation caused a decrease in the rate of heartbeat, and furthermore, the condensation of the water vapor produced a blurred image which interfered with the observations. In addition, the consequent reduction of the oxygen-tension level appeared to impair the respiratory function of the test animals, and this undoubtedly contributed in many cases to the recording of abnormally low levels of depression in rate of heartbeat.



Small double chambers were used and soon abandoned in favor of the following method and its variation. Although the double chamber could be used to project a normal and a test animal simultaneously, the technique involved in mounting the animals was too time consuming and possessed several of the disadvantages of the hanging-drop method.

The procedure, designated by us as the Mass Culture method, has demonstrated its merit. *Daphnia* from a standardized culture are divided into two groups. The first is placed in a one ounce bottle three-quarters filled with the drug solution which depressed the rate of heartbeat, and the second group is placed in a bottle similarly filled with a mixture of the depressant drug and digitalis or digitalis glucosides in varying proportions. At uniformly timed intervals the daphnia were rapidly removed, and the rate of heartbeat was determined in the depression of a ground glass slide. The protective activity of digitalis was manifested at that concentration where none or little depressant activity was observed in the mixture of depressant drug and digitalis. The level of depression caused by the depressant activity of the drug was obtained in the first group, and intermediate levels were obtained in certain concentrations of the drug and digitalis mixtures. Of course, the normal rate of heartbeat was determined at the beginning of the assay, and precautions were taken to maintain a constant temperature, usually 70 degrees F., at which temperature it has been our practice to culture the daphnia.

A variation of this method, designated as the Single Culture method, consists of the taking the normal heartbeats of many daphnia and placing them individually in the one ounce bottles similarly filled with the depressant drug and mixtures of the depressant drug and digitalis in various concentrations. In this way, the reactions of individual daphnia could be studied over long periods of time. The rate of heartbeat was counted as in the immediately preceding method. This Single Culture method can be employed after the injection of definite amounts of drug and drug-digitalis mixtures, and culture medium is used in their stead in the bottles. In this paper we shall report only those results which do not embody the microinjection technique.

IV. Results and Discussion

Yohimbine, because of its relatively low toxicity, was found to be the most suitable cardiac depressant for our purposes. Degree of

cardiac depression in daphnia is measured by the decrease in the number of heartbeats per minute. Three samples of digitalis were used: Powdered Digitalis Leaf (A), International Standard Digitalis (B), and a mixture of Digitalis glucosides (C).

Five daphnia (Table I) placed in 0.025 per cent. yohimbine solution, and having an average normal heartbeat of 267, at the end of two hours had an average heartbeat of only 190. On the other hand, the five daphnia which were placed in mixture of 0.025 per cent. yohimbine

TABLE II
EFFECT OF DIGITALIS LEAF PREPARATIONS AGAINST YOHIMBINE DEPRESSION

Normal Heart Beat				Normal Heart Beat				Normal Heart Beat			
1.	281	hrt.	bts. /min.	1.	283	hrt.	bts. /min.	1.	273	hrt.	bts. /min.
2.	287	"	" "	2.	292	"	" "	2.	285	"	" "
3.	300	"	" "	3.	292	"	" "	3.	277	"	" "
4.	283	"	" "	4.	269	"	" "	4.	269	"	" "
5.	290	"	" "	5.	274	"	" "	5.	277	"	" "
<hr/>				<hr/>				<hr/>			
Av. 288				Av. 282				Av. 276			
<hr/>											
One Hour											
.025% Yohimbine				.025% Yohimbine + .0745% Dig. (A) *				.025% Yohimbine + .0345% Dig. (A)			
1.	202	hrt.	bts. /min.	1.	198	hrt.	bts. /min.	1.	208	hrt.	bts. /min.
2.	263	"	" "	2.	214	"	" "	2.	205	"	" "
3.	225	"	" "	3.	222	"	" "	3.	217	"	" "
4.	214	"	" "	4.	243	"	" "	4.	191	"	" "
5.	209	"	" "	5.	273	"	" "	5.	209	"	" "
<hr/>				<hr/>				<hr/>			
Av. 223				Av. 190				Av. 206			
<hr/>											
Two Hours											
.025% Yohimbine				.025% Yohimbine + .0745% Dig. (A) *				.025% Yohimbine + .0345% Dig. (A)			
1.	181	hrt.	bts. /min.	1.	185	hrt.	bts. /min.	1.	180	hrt.	bts. /min.
2.	157	"	" "	2.	191	"	" "	2.	164	"	" "
3.	209	"	" "	3.	157	"	" "	3.	157	"	" "
4.	167	"	" "	4.	177	"	" "	4.	151	"	" "
5.	166	"	" "	5.	195	"	" "	5.	183	"	" "
<hr/>				<hr/>				<hr/>			
Av. 198				Av. 187				Av. 167			

* This sample of Digitalis was bought on the open market. It was more than two years old and was kept without precautions to prevent deterioration.

TABLE I
PROTECTIVE EFFECT OF DIGITALIS LEAF PREPARATIONS OVER A WIDE RANGE AGAINST YOHIMBINE DEPRESSION
Single Culture Method

Normal Heart Beat	Normal Heart Beat	Normal Heart Beat	Normal Heart Beat	Normal Heart Beat	Normal Heart Beat	Normal Heart Beat
1. 273 hrt. bts. /min.	1. 269 hrt bts. /min.	1. 275 hrt. bts. /min.	1. 292 hrt. bts. /min.	1. 277 hrt. bts. /min.	1. 277 hrt. bts. /min.	1. 295 hrt. b
2. 273 " " "	2. 257 " " "	2. 285 " " "	2. 269 " " "	2. 281 " " "	2. 269 " " "	2. 290 "
3. 253 " " "	3. 275 " " "	3. 271 " " "	3. 267 " " "	3. 273 " " "	3. 265 " " "	3. 288 "
4. 273 " " "	4. 250 " " "	4. 269 " " "	4. 269 " " "	4. 279 " " "	4. 277 " " "	4. 293 "
5. 300 " " "	5. 283 " " "	5. 277 " " "	5. 267 " " "	5. 285 " " "	5. 277 " " "	5. 295 "
Av. 274	Av. 267	Av. 275	Av. 273	Av. 278	Av. 273	Av. 292
<i>One Hour</i>						
Normal Heart Beat	.025% Yohimbine	0.1% Digitalis (A)	.025% Yohimbine + 0.1% Digitalis (A)	.025% Yohimbine + .05% Digitalis (A)	.025% Yohimbine + .025% Digitalis (A)	.025% Yohimbine + .0125% Digitalis (A)
1. 259 hrt. bts. /min.	1. 228 hrt. bts. /min.	1. 265 hrt. bts. /min.	1. 290 hrt. bts. /min.	1. 245 hrt. bts. /min.	1. 247 hrt. bts. /min.	1. 272 hrt. b
2. 269 " " "	2. 188 " " "	2. 271 " " "	2. 240 " " "	2. 279 " " "	2. 272 " " "	2. 273 "
3. 255 " " "	3. 173 " " "	3. 265 " " "	3. 285 " " "	3. 269 " " "	3. 261 " " "	3. 275 "
4. 269 " " "	4. 172 " " "	4. 283 " " "	4. 269 " " "	4. 257 " " "	4. 265 " " "	4. 281 "
5. 290 " " "	5. 172 " " "	5. 269 " " "	5. 240 " " "	5. 269 " " "	5. 273 " " "	5. 281 "
Av. 266	Av. 186	Av. 270	Av. 265	Av. 264	Av. 264	Av. 276
<i>Two Hours</i>						
Normal Heart Beat	.025% Yohimbine	0.1% Digitalis (A)	.025% Yohimbine + 0.1% Digitalis (A)	.025% Yohimbine + .05% Digitalis (A)	.025% Yohimbine + .025% Digitalis (A)	.025% Yohimbine + .0125% Digitalis (A)
1. 223 hrt. bts. /min.	1. 200 hrt. bts. /min.	1. 269 hrt. bts. /min.	1. 269 hrt. bts. /min.	1. 240 hrt. bts. /min.	1. 263 hrt. bts. /min.	1. 271 hrt. b
2. 248 " " "	2. 148 " " "	2. 273 " " "	2. 273 " " "	2. 285 " " "	2. 272 " " "	2. 273 "
3. 281 " " "	3. 250 " " "	3. 273 " " "	3. 273 " " "	3. 277 " " "	3. 277 " " "	3. 269 "
4. 288 " " "	4. 175 " " "	4. 277 " " "	4. 277 " " "	4. 277 " " "	4. 257 " " "	4. 275 "
5. 285 " " "	5. 180 " " "	5. 261 " " "	5. 261 " " "	5. 273 " " "	5. 269 " " "	5. 277 "
Av. 275	Av. 190	Av. 270	Av. 270	Av. 270	Av. 268	Av. 273

TABLE I

LIS LEAF PREPARATIONS OVER A WIDE RANGE AGAINST YOHIMBINE DEPRESSION

Single Culture Method

Normal Heart Beat	Normal Heart Beat	Normal Heart Beat	Normal Heart Beat
1. 292 hrt. bts. /min.	1. 277 hrt. bts. /min.	1. 277 hrt. bts. /min.	1. 295 hrt. bts. /min.
2. 269 " " "	2. 281 " " "	2. 269 " " "	2. 290 " " "
3. 267 " " "	3. 273 " " "	3. 265 " " "	3. 288 " " "
4. 269 " " "	4. 279 " " "	4. 277 " " "	4. 293 " " "
5. 267 " " "	5. 285 " " "	5. 277 " " "	5. 295 " " "
Av. 273	Av. 278	Av. 273	Av. 292

One Hour

.025% Yohimbine + 0.1% Digitalis (A)	.025% Yohimbine + .05% Digitalis (A)	.025% Yohimbine + .025% Digitalis (A)	.025% Yohimbine + .0125% Digitalis (A)
1. 290 hrt. bts. /min.	1. 245 hrt. bts. /min.	1. 247 hrt. bts. /min.	1. 272 hrt. bts. /min.
2. 240 " " "	2. 279 " " "	2. 272 " " "	2. 273 " " "
3. 285 " " "	3. 269 " " "	3. 261 " " "	3. 275 " " "
4. 269 " " "	4. 257 " " "	4. 265 " " "	4. 281 " " "
5. 240 " " "	5. 269 " " "	5. 273 " " "	5. 281 " " "
Av. 265	Av. 264	Av. 264	Av. 276

Two Hours

.025% Yohimbine + 0.1% Digitalis (A)	.025% Yohimbine + .05% Digitalis (A)	.025% Yohimbine + .025% Digitalis (A)	.025% Yohimbine + .0125% Digitalis (A)
1. 269 hrt. bts. /min.	1. 240 hrt. bts. /min.	1. 263 hrt. bts. /min.	1. 271 hrt. bts. /min.
2. 273 " " "	2. 285 " " "	2. 272 " " "	2. 273 " " "
3. 273 " " "	3. 277 " " "	3. 277 " " "	3. 269 " " "
4. 277 " " "	4. 277 " " "	4. 257 " " "	4. 275 " " "
5. 261 " " "	5. 273 " " "	5. 269 " " "	5. 277 " " "
Av. 270	Av. 270	Av. 268	Av. 273

Heart Beat	Normal Heart Beat			
s. /min.	1.	288	hrt. bts. /min.	
" "	2.	277	" "	" "
" "	3.	283	" "	" "
" "	4.	277	" "	" "
" "	5.	288	" "	" "
<hr/>				
Av. 280				

Yohimbine + Digitalis (A)	.025% Yohimbine + .00625% Digitalis (A)			
bts. /min.	1.	274	hrt. bts. /min.	
" "	2.	277	" "	" "
" "	3.	281	" "	" "
" "	4.	269	" "	" "
" "	5.	250	" "	" "
<hr/>				
Av. 270				

Yohimbine + Digitalis (A)	.025% Yohimbine + .00625% Digitalis (A)			
bts. /min.	1.	281	hrt. bts. /min.	
" "	2.	277	" "	" "
" "	3.	281	" "	" "
" "	4.	273	" "	" "
" "	5.	269	" "	" "
<hr/>				
Av. 276				

The first part of the report
 deals with the general
 situation of the country
 and the progress of the
 work during the year.

The second part of the report
 deals with the results of the
 work during the year.

The third part of the report
 deals with the results of the
 work during the year.

The fourth part of the report
 deals with the results of the
 work during the year.

The fifth part of the report
 deals with the results of the
 work during the year.

and 0.1 per cent. digitalis (A), and which had an average normal heartbeat of 273, at the end of two hours had an average heartbeat of 270, a decrease of only three beats. Five other animals which were kept in culture water instead of drug solution showed a variation in average heartbeat during the two hours which was essentially the same as in the yohimbine digitalis mixture. A solution of 0.1 per cent. digitalis (A) without yohimbine produced a depression of seven beats

TABLE III
EFFECT OF INTERNATIONAL STANDARD DIGITALIS AGAINST YOHIMBINE
DEPRESSION

Normal Heart Beat			Normal Heart Beat			Normal Heart Beat		
1.	270	hrt. bts. /min.	1.	290	hrt. bts. /min.	1.	277	hrt. bts. /min.
2.	295	" " "	2.	295	" " "	2.	285	" " "
3.	285	" " "	3.	285	" " "	3.	288	" " "
4.	272	" " "	4.	285	" " "	4.	279	" " "
5.	290	" " "	5.	300	" " "	5.	288	" " "
<hr/>			<hr/>			<hr/>		
Av. 282			Av. 278			Av. 283		
<hr/>								
One Hour								
.025% Yohimbine			.025% Yohimbine + .0745% Dig. (B) *			.025% Yohimbine + .035% Dig. (B)		
1.	225	hrt. bts. /min.	1.	212	hrt. bts. /min.	1.	231	hrt. bts. /min.
2.	247	" " "	2.	269	" " "	2.	235	" " "
3.	225	" " "	3.	176	" " "	3.	220	" " "
4.	212	" " "	4.	196	" " "	4.	192	" " "
5.	218	" " "	5.	222	" " "	5.	228	" " "
<hr/>			<hr/>			<hr/>		
Av. 225			Av. 215			Av. 223		
<hr/>								
Two Hours								
.025% Yohimbine			.025% Yohimbine + .0745% Dig. (B) *			.025% Yohimbine + .035% Dig. (B)		
1.	231	hrt. bts. /min.	1.	203	hrt. bts. /min.	1.	228	hrt. bts. /min.
2.	200	" " "	2.	214	" " "	2.	217	" " "
3.	189	" " "	3.	247	" " "	3.	214	" " "
4.	207	" " "	4.	219	" " "	4.	217	" " "
5.	180	" " "	5.	219	" " "	5.	218	" " "
<hr/>			<hr/>			<hr/>		
Av. 201			Av. 220			Av. 219		

* .0745% International Standard Digitalis solution is equivalent to one Cat Unit per 100 cc.

in the average heartbeat per minute, a depression which does not exceed the normal variation.

The protective action of digitalis (A) against 0.025 per cent. yohimbine was evident in concentrations as low as 0.0625 per cent.

A sample of the same preparation of digitalis, which was more than two years old and which had been kept with no precautions to prevent deterioration, exhibited no protective action. As a matter of fact, it seemed to augment rather than inhibit the action of yohimbine (Table II).

A smaller concentration of digitalis produced a corresponding decrease of protection (Table I). A concentration of 0.1 per cent. digitalis (A) allowed only a 5 per cent. depression; 0.00625 per cent. digitalis (A) allowed a depression of 11 per cent. This effect of graded dosage indicates the feasibility of using the protective action of digitalis as a basis for an assay.

More experiments by the Mass Culture method were carried out using 0.025 per cent. yohimbine and varying concentrations of digitalis glucosides. Ten animals, having an average normal heartbeat of 277, were placed in a 0.025 per cent. yohimbine solution. Ten other animals, whose average heartbeat was 270, were placed in a solution containing 0.025 per cent. yohimbine and 0.00167 per cent. digitalis glucosides. At the end of one hour the daphnia were removed one by one and their heartbeat counted and averaged. The first group had an average heartbeat of 216 and the group in the yohimbine-glucoside solution had dropped to 256 (Table IV). The time limit was set as one hour because the one hour and two hour readings were practically the same.

The protective action of digitalis glucosides (C) against 0.025 per cent. yohimbine was evident in a concentration as low as 0.00033 per cent.

Direct comparison can be made between samples (A) and (B) because they are both powdered leaf preparations. However, precise comparison of the protective action cannot be made between the powdered leaf preparations and digitalis (C), the glucosidal mixture, because of various other substances in the leaf preparations, which may directly or indirectly inhibit or augment the action of the cardiac glucosides present.

The results indicate that Digitalis (A) has a greater protective action than digitalis (B) at equivalent concentrations (Tables I and III). On the other hand digitalis (C) protects against 0.025 per

TABLE IV
 PROTECTIVE EFFECT OF DIGITALIS GLUCOSIDES AGAINST YOHIMBINE
 DEPRESSION
Mass Culture

Normal Heart Beat	Normal Heart Beat	Normal Heart Beat
286 Beats per min.	274 Beats per min.	286 Beats per min.
269 " " "	281 " " "	269 " " "
279 " " "	267 " " "	271 " " "
270 " " "	293 " " "	276 " " "
280 " " "	287 " " "	268 " " "
281 " " "	278 " " "	263 " " "
263 " " "	280 " " "	268 " " "
277 " " "	265 " " "	263 " " "
279 " " "	288 " " "	269 " " "
269 " " "	288 " " "	267 " " "
277 Av. beats per min.	280 Av. beats per min.	270 Av. beats per min.

One-half Hour

0.025% Yohimbine HCl	0.0011% Digitalis Glucosides	0.025% Yohimbine HCl + 0.00167% Digitalis Glucosides
246 Beats per min.	286 Beats per min.	247 Beats per min.
249 " " "	276 " " "	261 " " "
255 " " "	263 " " "	266 " " "
257 " " "	272 " " "	282 " " "
297 " " "	276 " " "	267 " " "
287 " " "	265 " " "	250 " " "
213 " " "	284 " " "	286 " " "
239 " " "	276 " " "	285 " " "
252 " " "	274 " " "	257 " " "
264 " " "	274 " " "	245 " " "
247 Av. beats per min.	275 Av. beats per min.	256 Av. beats per min.

One Hour

0.025% Yohimbine HCl	0.0011% Digitalis Glucosides	0.025% Yohimbine HCl + 0.00167% Digitalis Glucosides
215 Beats per min.	275 Beats per min.	278 Beats per min.
216 " " "	268 " " "	248 " " "
232 " " "	281 " " "	246 " " "
218 " " "	286 " " "	260 " " "
228 " " "	274 " " "	248 " " "
224 " " "	273 " " "	252 " " "
166 " " "	266 " " "	242 " " "
237 " " "	279 " " "	260 " " "
213 " " "	284 " " "	257 " " "
213 " " "	272 " " "	269 " " "
216 Av. beats per min.	278 Av. beats per min.	256 Av. beats per min.

cent. yohimbine in a concentration of about 1/20 of that of digitalis (A). But as mentioned above no really direct comparison can be made (Table V).

Before this non-toxic method of assay can become feasible, a number of factors producing variation in results must be further investigated and brought under control. The animals must be of the same age, sex, and of the same vitality. Animals which are anemic, small and generally sluggish will not produce the same results as healthier animals. The temperature must be constant.

Most important of all, a better understanding of standardization of the daphnia cultures must be obtained. At the present time, digitalis of a definite concentration does not always exhibit the same degree of protection in experiments in which all other factors apparently do not vary. Likewise, yohimbine in the concentrations used in these experiments does not always exhibit the same degree of depression. Thus, there are two variables which may be eliminated with the use of daphnia from more precisely controlled cultures.

V. Conclusions

1. Digitalis leaf exhibits protective action on the heart of *Daphnia magna* against depressants (e. g. yohimbine). The degree of protection appears to be directly proportional to the amount of digitalis.
2. Mixtures of digitalis glucosides exhibit the same protective action.
3. Further work is necessary to fully explore the feasibility of this method for assay purposes.

VI. Bibliography

- (1) Viehoveer, Arno: "Daphnia Propagation for Experimental Use." *A. J. Ph.*, 107, 3 (1935).
- (2) Viehoveer, A., and Cohen, I.: "Mechanism of Action of Aphrodisiacs." *A. J. Ph.*, 110, 6 (1938).
- (3) Viehoveer, Arno: "Daphnia, the Biological Agent." *J. A. Ph. A.*, 25, 12 (1936).
- (4) Viehoveer, A., and Cohen, I.: "Mechanism of Strychnine Action." *A. J. Ph.*, 109, 6 (1937).
- (5) Viehoveer, A., and Mikuriya, A. S.: "Effect of Digitoxin on the Heart of Daphnia." *J. A. Ph. A.*, 18, 1137 (1929). Abstract.
- (6) Viehoveer, A., and Tubis, M.: "Effect of Digitonin on the Heart of Daphnia." *Ibid.*
- (7) Viehoveer, A.: "Effect of Digitalis and Its Glucosides Upon the Chloroformed Daphnia Heart. Micro-Movies (1933).
- (8) Rowe, L. W.: "Further Evidence of the Strong and Variable Action of the U. S. P. XI Digitalis Standards." *J. A. Ph. A.*, 27, 10 (1938).

This work was carried on with the support and cooperation of Mr. J. Rosin, of Merck & Co.; the Sandoz Chemical Co., and John Wyeth & Bro., Inc.

TABLE V

AVERAGE RATE OF HEART BEAT INDICATING EFFECT OF GRADED DOSAGE
Averages Obtained From Twenty-five Experiments Using Digitalis (A)

Normal Heart Beat per Min.	282	281	275	282	278	279	274	281
Dosage	0.025% Yohimbine	0.025% Yoh. + 0.1% Digitalis	0.025% Yoh. + 0.075% Digitalis	0.025% Yoh. + 0.05% Digitalis	0.025% Yoh. + 0.025% Digitalis	0.025% Yoh. + 0.0125% Digitalis	0.025% Yoh. + 0.00625% Digitalis	0.025% Yoh. + 0.003125% Digitalis
Heart Beat after 2 Hours	209	266	266	256	244	258	244	198

Fig. 1

Averages Obtained From Fifteen Experiments Using Digitalis (B)

Normal Heart Beat per Min.	274	288	280	273	275	276
Dosage	0.025% Yohimbine	0.025% Yoh. + 0.1% Digitalis	0.025% Yoh. + 0.0745% Digitalis	0.025% Yoh. + 0.05% Digitalis	0.025% Yoh. + 0.0345% Digitalis	0.025% Yoh. + 0.025% Digitalis
Heart Beat after 2 Hours	196	239	220	223	204	180

Fig. 2

Averages Obtained From Thirteen Experiments Using Digitalis (C)

Normal Heart Beat per Min.	276	280	272	279	270	271	272
Dosage	0.025% Yohimbine	0.025% Yoh. + 0.0033% Dig. Gluc.	0.025% Yoh. + 0.00165% Dig. Gluc.	0.025% Yoh. + 0.0011% Dig. Gluc.	0.025% Yoh. + 0.00066% Dig. Gluc.	0.025% Yoh. + 0.00033% Dig. Gluc.	0.025% Yoh. + 0.000165% Dig. Gluc.
Heart Beat after 1 Hour	188	242	244	252	240	220	199

Fig. 3

EFFECT OF GRADED DOSAGE

Experiments Using Digitalis (A)

278	279	274	281
0.025% Yoh. + 0.025% Digitalis	0.025% Yoh. + 0.0125% Digitalis	0.025% Yoh. + 0.00625% Digitalis	0.025% Yoh. + 0.003125% Digitalis
244	258	244	198

Experiments Using Digitalis (B)

273	275	276
0.025% Yoh. + 0.05% Digitalis	0.025% Yoh. + 0.0345% Digitalis	0.025% Yoh. + 0.025% Digitalis
223	204	180

Experiments Using Digitalis (C)

269	270	271	272
0.025% Yoh. + 0.011% Dig. Gluc.	0.025% Yoh. + 0.00066% Dig. Gluc.	0.025% Yoh. + 0.00033% Dig. Gluc.	0.025% Yoh. + 0.000165% Dig. Gluc.
252	240	220	199

ABSTRACTS FROM AND REVIEWS OF THE LITERATURE OF THE SCIENCES SUPPORTING PUBLIC HEALTH

The Identity of an Antibacterial Factor in the Saliva of Certain Mammalia. S. L. Rosenthal, W. M. McNabb and R. C. Snyder. *J. A. D. A.* 26, 1859 (1939). Attempts to induce oral Vincent's infection in normal animals by inoculation have been unsuccessful. While localized infection has resulted in some cases it is not comparable with the disease in man. The lesions heal spontaneously and the infection is not transmitted to other animals in contact with the affected animals.

An investigation of this apparent immunity was begun by bacteriologic examination of the mouths of normal mammals including both domesticated and wild animals. Smears were notable for their low bacterial count in comparison with man's and for the complete absence of motile microorganisms. Fusiform bacilli or spirochetes similar to those found commonly in the human oral cavity were not found except in a few cases where some pathologic condition or an extremely dirty mouth existed.

Apparently, the mouths of healthy animals contained antibacterial substances since their habits and diets must bring them in contact with innumerable spirochetal forms yet not only is spirochetosis rare in these species but also the spirochete is seldom seen in healthy specimens.

In an attempt to identify such a substance saliva was collected from several animals of different species. The pH was found to be from 8.4 to 8.6. Each sample was subjected to the motility test which consists in bringing a definite quantity of the fluid under a dark field microscope with an equal quantity of saliva from a patient with Vincent's infection. The saliva contains actively motile spirochetes, vibrios and bacilli together with the natural organic debris. The antibacterial properties are measured by the time elapsing from the moment of contact of the solutions to cessation of movements of (1) spirochetes, (2) vibrios and (3) motile bacilli. Other workers have shown that loss of motility in *T. pallidum* entails loss of pathogenicity.

Contact with all normal animal saliva caused spirochetes and vibrios from a human mouth to lose motility instantly. Many instantly destroyed the motility of all bacilli as well but others failed to penetrate debris well or to stop the movement of all motile bacilli.

The pH of saliva from dogs which were household pets and suffering from Vincent's infection was found to be 6.7-7.6, about the range of that in normal man. These animals upon recovery were found to have a salivary pH of 8.6 and all had regained the inhibitory effect on the motile oral flora.

The constituent of normal animal saliva which possessed this antibacterial action was finally found to be sodium carbonate. Interesting was the additional finding that not in pH alone does the effect of sodium carbonate rest, since a solution of sodium hydroxide of similar pH is much less active.

Chemical tests were made on two dogs and five persons with acute Vincent's infections. Continuous topical applications of a molar solution of sodium carbonate were made to the dried gingival tissues for fifteen minutes. Home care was limited to hourly rinsing or topical applications with this solution. Resolution was rapid in all cases.

L. F. T.

Methyl Cellulose in Pharmacy. G. A. Bergy, *Amer. Prof. Pharm.* (V, 691 (1939)). Methyl cellulose dissolved in distilled water at normal temperatures forms a stable jelly. This property enables its use with simple procedures, to form mucilages resistant to extreme heat, cold and light and impervious to bacterial decomposition. Preservatives needed are merely to protect the other ingredients of the preparation.

Methyl cellulose is marketed as a coarse, dry fibrous mass which when dissolved in water produces a uniformly viscous product free from lumps.

A slight sediment may form on standing and may be removed by centrifugalization or decantation of the supernatant liquid.

Solutions of methyl cellulose are neutral in reaction, exhibiting excellent dispersing and emulsifying qualities. Emulsions made thus are strongly alkaline in hot or cold, and on cooling the clarity and thickness are improved. This is advantageous in preparations containing titanium dioxide, zinc stearate or carbon black for the shearing

effect of mechanical agitation advances and improves reduction of particle size.

The solutions of methyl cellulose are prepared by pouring 10 to 15 parts of boiling water over 1 part of the substance, stirring thoroughly until the mass swells and lumps dissolve, being careful not to incorporate too much air. Allow the solution to cool to approximately 60° C., stir, and cool completely, stirring again to make it homogeneous. Strain through unsized, unbleached muslin. The stock solution thus prepared may be diluted to any desired strength.

Solutions of methyl cellulose and products containing them are harmless to man and animals either internally or externally. Being resistant to marked changes in pH, the stomach's acidity and the intestines alkalinity do not decompose it.

The author suggests the use of methyl cellulose as a thickening or binding agent, and emulsifying or dispersing agent in pharmaceutical or dermatological preparations. Use the rule for tragacanth of seventy grains to the pint or 10 gms. to the liter for the strength to be employed. The use of methyl cellulose in dentifrices, skin creams, depilatories, hair grooms, cleansing creams and dermal creams with a typical formula for each is recommended.

M. H.

Sulfapyridine for Pneumonia Control. *Baltimore Health News*, 16, 172 (1939) through *Squibb Abstr. Bull.* 12, 1303 (1939). Sulfapyridine has been used extensively in the treatment of pneumonia in a number of hospitals in Baltimore with great reduction in mortality. In one large series of cases only 7 per cent. mortality occurred and in a series of 70 small children with measles and pneumonia none died. Sulfapyridine is more effective than antipneumococcic serum and therefore it will probably be more generally used in the treatment of pneumonia in the patients' home. It was recommended by the Pneumonia Control Committee of the Maryland State Board of Health as the chief therapeutic agent in the control of this disease. This does not mean that the use of serum should be discarded but rather that serum be used as a supplement especially in very severe cases and in those not promptly improved by sulfapyridine. It is suggested that sulfapyridine be employed also in the treatment of lobar and interstitial pneumonia. Careful check-up must be main-

tained against toxic reactions. It should be discontinued at once if the white blood cell count falls below 4, 500/cu. mm. with less than 50 per cent. polymorphonuclear leukocytes.

It should also be withdrawn if a hemoglobin fall of 30 per cent. or more is observed, if there is any clinical evidence of excessive hemolysis or if more than a very few red blood cells are found in the urine. Urinary precipitation of acetylsulfapyridine is less likely to occur if sodium bicarbonate and moderately large amounts of fluid are given.

L. F. T.

Deflagration with Sodium Peroxide as a Simple Analytical Process for the Determination of Halogen, Sulfur and Other Constituents in Organic Substances. R. Kraus. *Z. Anal. Chem.* 117, 243 (1939) through *Squibb Abstr. Bull.* 12, 1402 (1939). Deflagration of solid organic substances and certain liquids by quick and short heating with a large excess of Na_2O_2 in a covered nickel crucible by the method presented gives a quick and complete combustion with relatively little attack on the crucible. The determination of halogens, sulfur and phosphorus may then be accurately made. Viscous liquids are first mixed with MgO and liquids of high boiling point are absorbed in filter paper in the crucible before mixing with the Na_2O_2 . The method is limited for low percentage of Cl, S., etc., only by the "blank" of the Na_2O_2 . It is applicable to substances, such as chlorotoluidine, which are not readily decomposed by the Carius method. The method is particularly well suited for routine analysis.

L. F. T.

An Experimental Study of the Relation Between Concentration of Disinfectants and Time Required for Disinfection. F. W. Tilley. *J. of Bacteriology* (38, 499, 1939, No. 5). Different types of disinfectants act upon bacterial protoplasm in different ways. It can not be expected that the relation between concentration of disinfectant, and time required for disinfection can be expressed by a single formula applicable to all types. These factors do appear to be exponentially related over considerable ranges of concentrations and disinfection times in many disinfectants. In such cases the derived mathematical values may be useful in evaluating or predicting bactericidal efficiency.

The only variables allowable are: Concentration of disinfectant and time required for disinfection. The value n is known as the "concentration exponent" or "coefficient of dilution," denoting the degree to which dilution lessens the rate of disinfection. As an example, ethyl alcohol has a high value of n but loses its efficiency on dilution more rapidly than phenol which has an n value of about one half.

A modified Rideal-Walker technic was used to determine the bactericidal efficiencies of phenol, orthocresol, paracresol, orthobutylphenol, parabutylphenol, resorcinol, *n*-propylresorcinol, ethyl alcohol and *n*-butyl alcohol against *Staphylococcus aureus* and *Eberthella typhosa*. The following formula was employed to calculate n :

$$n = \frac{\log t_2 - \log t_1}{\log C_1 - \log C_2}$$

TABLE 4

Values of n determined at different temperatures

Disinfectant	Test Organism	10° C.	20° C.	30° C.	40° C.
Phenol	<i>E. typhosa</i>	7.9	7.5	5.9	5.2
	<i>S. aureus</i>	6.5	6.5	6.4	5.4
Orthocresol	<i>E. typhosa</i>	8.3	7.9	5.5	5.1
	<i>S. aureus</i>	7.8	7.6	8.1	6.8
Paracresol	<i>E. typhosa</i>	8.9	8.4	6.4	5.3
	<i>S. aureus</i>	8.2	8.7	9.4	7.0
O-Butylphenol	<i>E. typhosa</i>		9.2	7.2	
	<i>S. aureus</i>		8.5	9.1	
P-Butylphenol	<i>E. typhosa</i>		9.2	9.3	
	<i>S. aureus</i>		8.9	8.1	
Resorcinol	<i>E. typhosa</i>	4.6	5.1	5.2	
	<i>S. aureus</i>	4.4	5.0	6.2	
Ethyl alcohol	<i>E. typhosa</i>	12.7	11.4	8.8	
	<i>S. aureus</i>	11.4	11.1	8.5	
N Butyl alcohol ...	<i>E. typhosa</i>		11.9	10.3	
	<i>S. aureus</i>		11.8	10.1	

SOLID EXTRACTS

By Ivor Griffith, Ph. M., Sc. D., F. R. S. A.

Despite the form in which this information is presented it may be accepted as trustworthy and up-to-date. Original sources are not listed but they may be obtained upon request.

When an ungiving ash tray falls on a mahogany table it usually leaves its tell-tale mark. Such indentations, if they are not too deep, can usually be removed without difficulty by a paste of glycerin, water and iron oxide. Dip a hard felt pad in the paste and rub briskly back and forth over the scratched surface until the markings have disappeared. This procedure is especially adapted to removing shallow scratches. Deep gougings, however, require more specialized treatment. Many polishing materials used in mechanical methods utilize glycerin as an essential ingredient.

And speaking of glycerin, a commodity much more useful than most people know. Christmas decorators will do well to remember that soaking holly wreaths and similar fresh greenery in a 5 per cent. solution of glycerin in water keeps these Yuletide ornaments sprightlier and livelier than they might otherwise keep. Of course after the pre-soaking the hung-up wreaths, etc., should be allowed to drain thoroughly before being placed. This procedure is necessary only in the case of leaves and leaf or other live green ornaments used for interior decorations. Outside weather takes care of its own, but the warmth inside crisps too soon the cold loving evergreens.

If you want to transfer a favorite design or print on a news-sheet to paper, cloth, wood or other surfaces, this can usually be satisfactorily accomplished by employing the following simple solution of familiar ingredients:

Glycerin	1 1/2
Soap	4
Alcohol	10
Water	10

Wet thoroughly the print or drawing to be transferred. Remove excess liquid with a clean blotter and invert print onto the paper or cloth to which it is to be transferred. Lay a piece of paper over it and rub with a blunt article such as a knife handle.

The intimate chemical relationships existings between alkaloids, hormones and many dyestuffs are not always appreciated by pharmacologists. Many of the new synthetics also are first cousins to the coal tar dyestuffs. For instance, Sulfanilamide compounds, used in the war against disease on a score of fronts, promise to be useful in the scientific study of the undiseased tissues of animals and plants. One such has been found to be an effective "vital stain." Vital stains are dyes used on living tissues, which they color up and thus make easier to examine under the microscope. The sulfanilamide compound used has been long since known under the trade name of Neoprontosil, and it tinges the cells of plants and insects red. The discovery was made incidentally, while investigating possible effects on virus diseases.

Any new use for castor oil to divert it from the use that you think of when you hear the name is news. Castor oil is being used to make paint, replacing the tung oil largely imported from China and now difficult to get on account of the Japanese invasion. As pressed from the castor bean, the oil will not dry properly. So it is necessary to dehydroxylate it, snatch out some hydrogen and oxygen (water, if you must know) atoms from its molecule, which makes it a drying, odorless, non-yellowing, oil comparable with tung oil. It is a 9-11 octadecadieneic acid, if you talk chemistry; Dehydrol, if you talk paint manufacture. As a partner to the dehydrated castor oil, there is phenol resin, one of the most familiar synthetic plastics made from carbolic acid (phenol) and formaldehyde, modified into a pale, extra hard resin for making quick drying, water resistant varnishes, enamels and undercoats.

Wounds in plants are caused to heal by an acid which has been isolated in crystalline form by Drs. James English, Jr., James Bonner and A. J. Haagen-Smit of the California Institute of Technology,

and for which they propose the name "traumatic acid." The substance has the chemical formula $C_{12}H_{22}O_4$, and is identical with the organic acid, 1-decene-1, 10-dicarboxylic acid. The three experimenters, using a solution of this acid synthetically prepared, induced rapid formation of healing tissue on the cut surfaces of potato tubers. The discovery of the chemical nature of the wound hormone is announced briefly in "Science," with the statement that a more detailed report will be published later, when its application to animal wounds may be discussed.

A chemical compound that may prove as good a remedy against infectious diseases as sulfanilamide, with less toxic effects, is announced by the U. S. Public Health Service.

The compound, prepared by Drs. Hugo Bauer and Sanford M. Rosenthal, at the National Institute of Health, contains phosphorus instead of sulfur, and is different in other ways from sulfanilamide. Three such compounds have been prepared, of which one, bis (4-dimethylaminophenyl) phosphinous acid, checked streptococcus infections in mice and had a low toxicity.

No human trials of these chemicals have been made yet, nor will they be, Dr. Rosenthal said, before more extensive laboratory investigations.

The object of the research, in which compounds with arsenic substituting for the sulfur of sulfanilamide were also made, is to find chemicals which either are better than sulfanilamide or are effective against germ infections which sulfanilamide does not check.

Listerine has listened to the law! Thousands who have hearkened to the outrageous claims made by its manufacturers and who have sought through it some surcease from the dandruff distribution will find little comfort in the fact that the Lambert Pharmacal Company have entered into a stipulation with the Federal Trade Commission to cease making the following unwarranted advertising claims for this product:

(1) That all dandruff is due to an infection with *Pityrosporon ovale* or any other organism;

- (2) That dandruff necessarily is a germ disease;
- (3) That the dandruff germ has been isolated or identified;
- (4) That the presence of *Pityrosporon ovale* necessarily means dandruff or that with its destruction dandruff disappears;
- (5) That dandruff is necessarily infectious, contagious or "catching" or is in all instances passed from one person to another, or
- (6) That any of the foregoing assertions is a "scientific fact" or a "fact definitely established by scientists."

It was also stipulated that the company stop representing that the product either cures or permanently relieves dandruff; that the product "kills the dandruff germ," "attacks the cause of dandruff," or penetrates infected hair follicles, or "annihilates" the dandruff germ.

Finally, the company will no longer advertise: that the product has "marked curative properties due to certain ingredients in a unique combination shared by no other antiseptic"; that ordinary remedies "aren't even antiseptic," are "smelly," affect only surface symptoms, or merely remove surface symptoms temporarily, or that competitive products are obviously inferior to "Listerine Antiseptic" as a remedy for dandruff—when such are not the facts.

But despite these capitulations there still seems to be a demand for Listerine if only because of its name.

BOOK REVIEWS

Done by persons, unafraid to upbraid, but perfectly willing to give praise where praise is really due.

A Textbook of Materia Medica, Pharmacology and Therapeutics. By Harold N. Wright, M.S., Ph.D., Associate Professor of Pharmacology, University of Minnesota; and Mildred Montag, R.N., M.A., Instructor in Nursing Arts, St. Luke's Hospital, New York City. 566 pages with 83 illustrations. Philadelphia and London: W. B. Saunders Company, 1939. Cloth, \$2.75.

This book is obviously intended to be used by pupil nurses, as is shown both by the subjects covered and the method of presentation. The first seventy-five pages are given over to an outline of some points of elementary pharmacy which are of interest to the nursing profession; then follows about fifteen pages chiefly devoted to consideration of various methods of administering drugs. This leaves about four hundred pages to be devoted to the subject of pharmacology. Even in this latter part of the book the authors have in mind always their intended audience and continuously emphasize those pharmacological facts which especially concern the nursing profession, with a compensatory skimping of many of the phases which are of interest to pharmacists.

The pharmacological portion of the book displays a pleasing clarity of style which is enhanced by a number of illustrative diagrams but is unfortunately marred by weird typographical arrangement. Here and there the effort at simplicity has lured the author into some generalizations which are not strictly accurate—such as the implication that all germicides act by precipitating protein—but anyone who can read the book unperturbed by the inconsistencies in the use of black-face type and in the paragraphing should obtain an understanding of the modern ideas of drug action. The drugs considered are not limited to those official but include a considerable number of the more important of the proprietary synthetics.

H. C. Wood.

INDEX TO VOLUME 111 OF THE AMERICAN JOURNAL OF PHARMACY

AUTHORS, 1939

- (A)—Abstract
(B)—Book Review
(E)—Editorial
(S)—Solid Extract

	PAGE		PAGE
Beuhrer, T. E.—		Couch, James F.—	
The Chemical Composition of		Hydrogen Ion Concentration and	
Rayless Goldenrod	105	Cyanogenesis in Sorghum ..	55
Billinger, R. D.—		Crowder, J. A.—	
America's Oldest Pharmacy ...	234	The Chemical Composition of	
Briese, Reinhold R.—		Rayless Goldenrod	105
Hydrogen Ion Concentration		Dickhart, W. H.—	
and Cyanogenesis in Sorghum	55	Chemical and Physical Prop-	
Brillhart, Russell E.—		erties of Argan Oil	293
Efficiency of Bactericidal Agents		Eisenberg, H.—	
in Different Ointment Bases.	430	On the Nature of the Precipi-	
Christiansen, W. G.—		tates Obtained by Adding	
Oxalate Formation in Ascorbic		Sodium Silicate to Magnesium	
Acid Solutions	347	Sulfate	400
Clark, W. C.—		Fairlamb, Philip M.—	
Antiseptic Value of Certain		A Survey of the Viscosities of	
Phenolic Ointments	228	Medicinal Mineral Oils	139
Cohen, Isadore—		Fulton, Charles C.—	
The Evaluation of the Activity		Alkaloids and Their Reagents.	184
of Powders of Veratrum		Gershenfeld, Louis—	
Viride by the Daphnia Method	426	Efficiency of Bactericidal Agents	
Physiological Evaluation of		in Different Ointment Bases.	430
Veratrum Viride and V.		Miscibility of Water Immiscible	
Album I. Toxicity	86	Disinfectants	314
Cook, E. Fullerton—		Orthotolidine and Orthotoluidine	
The New Pharmacopœia (E) ..	310	Tests for Occult Blood	17

	PAGE		PAGE
Preservatives for Nasal Jellies and Ophthalmic Solutions ...	385	Harrisson, J. W. E.— Experimental Pharmacology and Materia Medica (B)	419
Preservatives for Preparations Containing Gelatin	277	Jurist, A. E.— Oxalate Formation in Ascorbic Acid Solutions	347
Greene, L. Wilson— Bitter Almond Oil, Chemical Microscopy	10	Kleckner, L. G.— Industrial Solvents (B)	460
Griffith, Ivor— Biologic Cleanliness (E)	378	Konnerth, R. A.— Ionizable Silver or Strong Sil- ver Protein in Mild Silver Protein	65
Breaking the Fast (E)	342	Kramer, John E.— American Way of Life (B) ..	223
Crystal Gazing	321	Lobsnows (B)	376
Drugstore of the Future (E) ..	2	Landeem, Hazel E.— Why a Hospital Pharmacy? ...	359
Fundamentals of Pharmacy, Theoretical and Practical (B)	459	Last, Jules H.— Watermelon Diuresis, a Study of	68
In the Interest of Pharmacy (E)	136	Levin, M. B.— On the Nature of the Precipi- tates Obtained by Adding Sodium Silicate to Magnesium Sulfate	400
Origin of Habit (E)	226	Little, Ernest— An Important Responsibility of the Pharmaceutical Educator.	48
A Negligible Trace (E)	422	Mason, C. M.— The Chemical Composition of Rayless Goldenrod	105
The 1940 Pharmacopoeial Con- vention (E)	182	McDonnell, Joseph F., Jr.— A Survey of the Viscosities of Medicinal Mineral Oils	139
Petrillized (E)	84		
re-Search (E)	274		
Solid Extracts, 40, 131, 175, 301, 371, 416,	456		
William Procter, Jr. (E)	44		
Harding, T. Swann— Alcohol, Health, Longevity and Offspring	351		
American Medicine Mobilizes (B)	306		
Have a Smoke?	443		
The Remedies Are Not What They Used to Be	245		
Tragic Neglect of Scientific Methods	19		
What Is a "Cold"—and Then What?	143		

	PAGE		PAGE
McFarlan, Ronald L.—		Rancidity in Edible Fats (B) ..	305
Physics in Pharmacy	113	Standard Chemical and Technical Dictionary (B)	340
Olsen, Paul—		Treatise on Pharmacy (B) ...	178
Foreign Markets for American Medicinal Products (B)	305	Tomkin, David—	
Osol, Arthur—		Preservatives for Nasal Jellies and Ophthalmic Solutions ..	385
Assay of a Sample of Opium Ninety Years Old	380	Van Deripe, F. N.—	
Perlstein, David—		Ionizable Silver or Strong Silver Protein in Mild Silver Protein	65
Preservatives for Preparations Containing Gelatin	277	Viehoever, Arno—	
Pfeiffer, Carl—		A New Approach to the Standardization of Digitalis	466
Watermelon, Diuresis, a Study of	68	Marihuana, America's New Drug Problem (B)	178
Podolsky, Edward—		Physiological Evaluation of Veratrum Viride and V. Album I. Toxicity	86
The Conquest of Hemophilia..	288	Vitamins and Hormones (B) ..	179
Roby, Charles—		Watson, E. M.—	
Watermelon Diuresis, a Study of	68	Some Aspects of Fifty Years Progress in Pharmaceutical and Related Sciences	161
Roseman, R.—		Witlin, Bernard—	
On the Nature of the Precipitates Obtained by Adding Sodium Silicate to Magnesium Sulfate	400	Miscibility of Water Immiscible Coal-Tar Disinfectants	314
Sokoloff, N. H.—		Wood, H. C.—	
A New Approach to the Standardization of Digitalis	466	Assay of a Sample of Opium Ninety Years Old	380
Taransky, A. A.—		Manual of Toxicology (B) ...	419
A New Approach to the Standardization of Digitalis	466	Medical Jurisprudence and Toxicology (B)	272
Tice, L. F.—		Medullary Stimulants (A)	260
Biological Products (B)	460	A Textbook of Materia Medica, Pharmacology and Therapeutics (B)	484
The Extra Pharmacopeia (B). ..	82		
A Further Contribution on the Use of Gelatin in Emulsion Products	4		

TITLE AND SUBJECT INDEX FOR 1939

(A)—Abstract
 (B)—Book Review
 (E)—Editorial
 (S)—Solid Extract

	PAGE		PAGE
Acetone, Determination of (A) ..	365	Anaemia (E)	423
Acidosis	149	Anaesthesia	204
Acriflavine as a Preservative	387	Analeptic (A)	260
Adjuvant Medicinals (S)	456	Aneurin	116
Adrenalin Solutions (A)	170	Antibacterial Factor in the Saliva of Certain Mammalia (A)	475
Aerosol (E)	274	Anti-Fogging Compositions	18
Alcohol	279	Antimalarials, Synthetic (A) ...	192
Alcohol, Health, Longevity and Offspring	351	Antioxidants for Castor Oil (A). .	410
Alcohol Intoxication (A)	126	Antiseptic Value of Certain Phenolic Ointments	228
Algin (A)	294	Aplopappus Discoideus	108
Alkaloids and Their Reagents ...	184	Aplopappus Hartwegi	105
Alkalosis	144	Argon	161
Almond Oil, Bitter Chemical Microscopy	10	Argan Oil, Chemical and Physical Properties of	293
Aloe (E)	275	Ascorbic Acid (S)	372
Alpha and Beta-Eucaine	204	Ascorbic Acid Solutions	347
Ameba Behavior (S)	375	Aspirin, Gastric Irritation (A)..	414
America's Oldest Pharmacy	234	Assay of a Sample of Opium Ninety Years Old	380
American Medicine Mobilizes (B)	306	Atebrine (S)	270
American Pharmaceutical Asso- ciation, 1939 Convention Ab- stracts	364	Atmospheric Contamination (S). .	176
American Pharmaceutical Asso- ciation, Origin of (E)	45	Auramine (A)	16
American Way of Life (B)	223	Authentic Germs, Home for (S). .	132
Ammoniated Mercury as a Bac- tericidal Agent	430	Avertin (A)	172
Amyl Phenol, Para	230	Avocado Pear Oil (A)	77
Amylases (A)	128	Azochloramid as a Preservative. .	386
Amylocaine	205	Bactericidal Agents in Ointment Bases	430

PAGE	PAGE
Bactericidal Effects of Ultraviolet Radiation Produced by Low Pressure Mercury Vapor Lamps (A)	408
Bacteriology Review	212
Bathing (E)	378
Benzaldehyde	10
Benzoic Acid as a Preservative..	386
Benzyl Benzoate (S)	371
Betanaphthol	278
Betanaphthol as a Preservative..	386
Betanaphthol, Phenol Coefficient of	317
Bezoar Stones	332
Bichloride of Mercury as a Bactericidal Agent	430
Bigger and Better Bubbles (A).	346
Biochemistry Review	208
Biologic Cleanliness (E)	378
Biological Products (B)	460
Bird Muscle to Control Bleeding.	289
Birds' Number Sense (S)	304
Bismark Brown (A)	16
Bitter Apple (E)	275
Blueberry Juice (S)	268
Boneset (E)	275
Book Reviews, 82, 178, 223, 272, 305, 340, 376, 484	
Boric Acid as a Preservative....	386
Breaking the Fast (E)	342
Bubbles (A)	346
Buffer Action	167
Buffered Local Anesthetic Solutions (A)	220
Burton, Robert (S)	132
Bush Disease (E)	424
Butoben as a Preservative	387
Butyl Alcohol (A)	479
Butyl Hydroxy Benzoate	279
Butylphenol, Ortho (A)	479
Butylphenol, Para (A)	479
Butyn	205
Cadmium, Quinaldinic Acid as a Reagent for (A)	74
Calcium Acetylsalicylate (A) ...	318
Calcium Gluconate, Determination (A)	37
Calcium Level During Pregnancy (A)	125
Calcium Salts' Effect Upon Lactose Utilization (A)	295
Cancer-Producing Substances ...	212
Cantharidin (A)	367
Carbon Dioxide as an Analeptic (A)	261
Carbon Monoxide in Medicinal Oxygen (A)	365
Cardiazol (A)	262
Casein and Zein as New Soap Additives (A)	183
Castor Oil, New Products of (S)	417
Castor Oil Antioxidants (A) ...	410
Castor Oil Cocktail (S)	417
Castor Oil in Paint (S)	481
Cetyl Alcohol (A)	76
Chamomile (E)	275
Chemotherapy	206
Child Geniuses Succeed (S)	175
Chlorthymol	230
Chloramine	279
Chlorobutanol	278
Chlorobutanol as a Preservative..	387
Chlorothymol	278
Chlor-Symmetrical Xylenol, Phenol Coefficient of	317
Chlorthymol (Sat. Sol.) as a Preservative	386
Chlorthymol, Phenol Coefficient of	317
Chrysoidine (A)	16

	PAGE		PAGE
Citrate Ricinoleate Agar (A) ...	34	Creosote, Phenol Coefficient of..	317
Clauden	292	Crystal Gazing	321
Coagulen	291	Cube Root (E)	274
Coal-Tar Disinfectants	314	Culture Collection (S)	132
Cobalt Deficiency (E)	424	Cyanogenesis in Sorghum	55
Cod Liver Oil Emulsion, Oil Con- tent (A)	450	Cyanogenetic Plants	55
Coffee Stirring (S)	176	Cyclopropane (A)	172
Coffee Tars Cause Cancer (S)..	373	Damiana Leaf Oil	108
Colchicine (S)	371	Dandelion (S)	372
Colchicine in a New Role (A)..	369	Dandruff (S)	482
"Cold"	143	Daphnia Magna	91, 466
"Cold," What Is a—and Then What?	139	Daphnia Method of Evaluating Veratrum Viride	426
Coliform Analysis (A)	34	Deflagration With Sodium Perox- ide (A)	478
Collagen (S)	176	Depilatory Powder	243
Colloidal Bunk	324	Dermatologic Paste (A)	364
Colloidal Preparations in Medi- cines	168	Derris (E)	274
Color Change in Organic Solu- tions (A)	35	Design Transfer (S)	480
Color Reaction for Sulphur (A). 336		Detergents (E)	274
Colored Eggs and Rabbits (S)..	175	Digitalis (E)	275
Colorimetry	116	Digitalis Standardization, New Approach to	466
Copper Deficiency (E)	424	Disinfectants and Time Required for Disinfection (A)	478
Copper "Licks" (E)	424	Distilled Water, Colloidal Im- purities in (A)	276
Copper, Quinaldinic Acid as a Reagent for (A)	74	Divinyl Ether	203
Copper Sulfate	279	Doctrine of Signatures	237
Copper Sulfate as a Preservative 386		Drene (E)	274
Coramin (A)	263	Drug Therapy in Coronary Dis- ease (A)	73
Coronary Disease (A)	73	Drugs, Choice of (A)	129
Cosmetic Label Restrictions (S). 374		Drugstore of the Future (E) ...	2
Costal Sickness (E)	424	Education in Pharmacy (E) ...	136
Cow Powder	243	Educator, Responsibility of	48
Cresol	278	Efficiency of Bactericidal Agents in Different Ointment Bases ..	430
Cresol, Phenol Coefficient of....	317	Electrotherapy	164
Cresol, Ortho (A)	479		
Cresol, Para (A)	479		

PAGE	PAGE
Emulsification and Chemical Reaction (A)	78
Emulsions Using Gelatin	4
Equisetum (E)	275
Estrogen (A)	298
Ether Explosions (A)	81
Ethyl Alcohol (A)	479
Ethyl Hydroxy Benzoate	279
Ethyl Hydroxy Benzoate of Sodium as a Preservative	386
Evipan Sodium (A)	172
Examination of Pharmacists for the Medical Administrative Corp, Regular Army (A)	320
Experimental Pharmacology and Materia Medica (B)	419
Eye Water	243
Fat Determination in Small Amounts (A)	454
Feather Scales (S)	131
Fever and Ague Powder	242
Filtration Sterilization (A)	127
Flash-Pasteurization Processes (S)	269
Flash-Pasteurization Processes (E)	275
Flower Preservatives (S)	271
Fluorescence	115
Foreign Markets for American Medicinal Products (B)	305
Formaldehyde as a Preservative.	386
Formaldehyde in the Presence of Methenamine, Determination of (A)	219
Fruit Juices (E)	275
Fundamentals of Pharmacy, Theoretical and Practical (B)	459
Gallstones	331
Gardinol (E)	274
Gastric Irritation From Aspirin (A)	414
Gelatin Preparations, Preservatives for	277
Gelatin in Emulsion Products	4
"Gingin" Disease (E)	424
Glass Is a Heat Trap (S)	40
Glycerine (S)	480
Glycerin Colors (A)	16
Glycerine-Toughened Soap Solutions (A)	346
Goiter (S)	177
Good Old Days (S)	42
Grippe	144
Gout	334
Guaiacol, Phenol Coefficient of..	317
Guava Juice (S)	268
Habit, Origin of (E)	226
Hair Clues in Crime Detection (S)	416
Harlem Oil	242
Have a Smoke?	443
Hay Fever Absent in Japan (S).	373
Hay-Fever Pollen (S)	457
Hegari	156
Helium	161, 172
Hemophilia, Conquest of	288
Henry VIII Fancied Pharmacy (A)	384
Heparin (A)	337
Hexylresorcinol (A)	124
Hexylresorcinol as a Preservative.	386
Hexylresorcinol in Pinworm Infection (A)	452
Hexylresorcinol, Phenol Coefficient of	317
History of Opium Assay	381
Honey (S)	456
Hormone, Unrecognized (S) ...	302
Horse Nettle (E)	275
Horse Powder	242

	PAGE		PAGE
Horsetail (E)	275	Liquor	351
Hospital Pharmacy, Why a	359	Liverwort (E)	275
Hospital Pharmacy Standards ..	361	Lobscows (B)	376
Hospital Statistics	360	Lung Extract to Control Bleeding	292
Hydrogen Peroxide38, 279, 387		Magnesium Citrate Solution	319
Hydrogen Peroxide for Pharma- ceutical Purposes, a Note on the Stabilization of (A)	338	Magnesium Sulphate and Sodium Silicate Precipitates	400
Hydrogen Ion Concentration and Cyanogenesis in Sorghum	55	Magnesium Sulfate in the Meas- urement of Circulation Time (A)	171
Hysterical Drops	242	Malachite Green (A)	16
Impatiens Plant (E)	275	Malachite Green as a Preserva- tive	387
Industrial Solvents (B)	460	Malaria, Test for (A)	172
Influenza	143	Malaria, Warning Against Treatment of Induced With Mapharsen and Tryparsamide (A)	335
Insecticides (E)	274	Mammary Carcinoma (A)	298
Insulin, Significance of the Zinc Content of Commercial Samples of (A)	126	Manual of Toxicology (B)	419
Iodides Incompatible with Strych- nine (A)	266	Marihuana, America's New Drug Problem (B)	178
Iodine as a Preservative	387	Medical Jurisprudence and Toxi- cology (B)	272
Iodocholeate (A)	366	Medicinals, Manufacture of (E).	138
Ionizable Silver	65	Medinal (A)	172
Ionizable Silver or Strong Silver Protein in Mild Silver Protcin.	65	Medullary Stimulants (A)	260
Irium (E)	274	Meigs, Dr. Charles, Biography..	380
Iso-valeric Acid (S)	371	Mercarbolid as a Preservative..	386
Ivy Poisoning, Treatment and Prevention (A)	412	Mercuric Chloride as a Pre- servative	386
Jenner, Edward (S)	133	Mercurophen as a Preservative..	386
Johnson Grass	151	Merthiolate as a Preservative ...	386
Kidney Stones	333	Metaphen as a Preservative	386
Krypton	161	Methyl Cellulose in Soap Making (A)	173
Leeches (S)	134	Methyl Hydroxy Benzoate	279
Lettuce (E)	275	Methyl Hydroxy Benzoate of Sodium as a Preservative	386
Limnology (S)	458	Methyl Violet (A)	16
Liquid Petrolatum, Prevention of Carotene Absorption (A)	451		
Listerine (S)	482		
Lithium Craze in Medicine	333		

PAGE	PAGE
Methyl Violet as a Preservative. 386	Ophthalmic Solution Preservatives 385
Methylene Blue (A) 16	Opium, Assay of Ninety Year Old Sample 380
Metrazol (A) 262	Organic Chemistry Review 202
Milkweed (S) 372	Orthophenylphenol, Phenol Coeffi- cient of 317
Milkweed Enzyme (S) 457	Orthotolidine and Orthotoluidine Tests for Occult Blood. 17
Mineral Oils 139	Oscillograph 166
Miscibility of Water Immiscible Coal-Tar Disinfectants 314	Oxalate Formation in Ascorbic Acid Solutions 347
Morphine Synthesis Unsuccessful (S) 269	Oxygen, Carbon Monoxide in Medicinal (A) 365
Morphinometric Assay, History of 380	Oxyquinoline Sulfate 278, 386
Mosquito (A) 78	Parachlormetacresol (A) 124
Mullein (E) 275	Parachlormetacresol 278, 317, 336
Musk, Artificial and Synthetic (S) 301	Parachlormetaxyleneol 230
Mycose (S) 41	Parachlormetaxyleneol 278
Nasal Hygiene (A) 170	Paraldehyde (A) 172
Nasal Jelly Preservatives 385	Parathyroid, Pharmacognosy of (A) 365
Nectars (E) 275	Passion Fruit Juice (S) 268
Negligible Trace, A (E) 422	Pastes for Dermatologic Use (A) 364
Nembutal (A) 172	Penetrating Agents (E) 274
Neon 161	Pentothal Sodium (A) 172
Neoprontosil (S) 481	Percaïne (A) 172, 205
New Approach to the Standard- ization of Digitalis 466	Petrillized (E) 84
Nicotine (S) 304	Petrolatum, Liquid 139
Nitrophenylhydrazine, Para 13	Petrolatums, Natural and Arti- ficial (A) 338
Nose Drops 147	pH 167
Novacaine (A) 172	pH Effect on the Bactericidal Properties of Commercial Anti- septics (A) 218
Nupercaine (A) 172	Pharmacognosy of the Parathy- roid (A) 365
Occult Blood, Tests for 17	Pharmacopœial Convention 1940 (E) 182
Occupational Diseases of the Pharmacist (A) 265	Pharmacy, In the Interest of (E) 136
Oil Content in Emulsion of Cod Liver Oil (A) 450	
Ointment Bases, Bactericidal Agents in 430	
Operations (S) 134	

	PAGE		PAGE
Pharmacy in Public Health (E) .	462	Position of Pharmacy in Public Health (E) .	462
Pharmagel	4, 277	Posterior Pituitary Extract (A) .	220
Pharmacopœia, the Extra (B) ..	82	Potassium or Sodium Sulfo- cyanate (A)	367
Pharmacopœia, the New (E) . . .	310	Potentilla Anserina (S)	372
Pharmacopœial Supplements (E)	310	Poultry Preparation (S)	40
Phenol	278, 479	Precipitates Obtained by Adding Sodium Silicate to Magnesium Sulphate	400
Phenol as a Bactericidal Agent..	430	Preservatives for Nasal Jellies and Ophthalmic Solutions . . .	385
Phenolic Ointments	228	Preservatives for Preparations Containing Gelatin	277
Phenylhydrazine	12	Procaine (A)	172, 205
Phenyl Hydroxy Mercuric Chlor- ide as a Preservative	387	Procaine Hydrochloride Solutions for Surgical Use (A)	36
Phenyl Mercuric Acetate as a Preservative	387	Procter, William, Jr. (E)	44
Phenylmercuric Nitrate (A), 124, 336, 387		Progress in Pharmaceutical and Related Sciences	161
Paraphenyl Phenol	230	Propyl Hydroxy Benzoate	279
Philadelphia College of Pharmacy and Science, 117th Commence- ment	255	Propyl Hydroxy Benzoate of Sodium as a Preservative . . .	386
Phosphinous Acid, Bis (4-Dimeth- ylaminophenyl) (S)	482	Prostigmin in the Treatment of Deafness (A)	299
Photoelectricity	118	Prunus Serotina	62
Photometer	115	Public Health, Position of Phar- macy in (E)	462
Physics in Pharmacy	113	Purpura	288
Picrotoxin (A)	261	Pylam Basic Black (A)	16
Pill, the Story of the (A)	263	Pylam Scarlet (A)	16
Pinworm Infection (A)	451	Pyrethrum (E)	274
Plaintain (E)	275	Pyridine	110
Plasmochin (S)	270	Quinaldinic Acid as a Reagent for Copper, Zinc and Cadmium (A)	74
Pneumococcic Infection (A) . . .	126	Quinine Ethyl Carbonate	279
Pneumococcic Pneumonia (A) ..	79	Quinine Synthesis Unsuccessful (S)	269
Pneumonia Death Rate in Nor- way (S)	374	Radioactive Iron (A)	217
Poison (S)	42	Radioactive Substances	162
Poisons (E)	138		
Pokeweed (S)	372		
Polarization	117		
Poliomyelitis Virus in the Stools of Healthy Contacts (A)	453		

	PAGE		PAGE
Radio-Sodium	162	Snake Venom to Control Bleeding	288
Radium	161	Soap Additives (A)	183
Radon	161	Soap Making (A)	173
Ragweed (S)	373	Soaps, Pharmacology of (A) ...	366
Rancidity in Edible Fats (B) ...	305	Soap Substitutes in Dentifrices (A)	409
Rau Drug Store	234	Sobering-up Treatment (S)	304
Rayless Goldenrod	105	Sodium Acid Sulfite	14
Refrigerated Blood (S)	418	Sodium Amytal (A)	172
Remedies Are Not What They Used to Be	245	Sodium Benzoate	278
re-Search (E)	274	Sodium Benzoate as a Preserva- tive	387
Resorcinol (A)	479	Sodium Perborate (A)	38
Resorcinol, Phenol Coefficient of.	317	Sodium or Potassium Sulfo- cyanate (A)	367
Responsibility of the Pharma- ceutical Educator	48	Sodium Salicylate	278
Rhubarb Juice (S)	268	Sodium Salicylate as a Preserva- tive	387
Roborant (A)	260	Sodium Silicate and Magnesium Sulphate Precipitates	400
Rotenone (E)	274	Sodium Silico Fluoride as a Pre- servative	386
Saccharin (S)	133	Sodium Thioethamyl (A)	172
Sage (E)	275	Soft Paraffins, Natural and Arti- ficial (A)	338
Salicyl Anilide (Sat. Sol.) as a Preservative	387	Solid Extracts, 40, 131, 175, 268, 301, 371, 480	
Salicylic Acid, Phenol Coefficient of	317	Sorghum	55
Santomer D	232	Sorghum Halepense	61, 151
Santophen "7"	230	Sorghum Vulgare	61
Saponated Solution of Cresol (A)	364	Sorrel (S)	372
Scratch Remover (S)	480	Sound Waves—A New Tool for Food Manufacturers (A)	296
Sebacic Acid (S)	417	Spectrophotometric Assay	166
Seconal (A)	172	Spectrophotometric Examination of Blood From Animals Re- ceiving Sulfanilamide (A) ...	264
Selenium Poisoning (S)	371	Spectroscopy	113
Semicarbazide	13	Spending Statistics (S)	456
Senyiga	143	Spinach (S)	372
Serums for Bleeding	290		
Silver Protein	65		
Simes, Samuel, Biography	381		
Skunk Cabbage (S)	372		
Smallpox (S)	133		
Smoking Habits and Statistics ..	443		

	PAGE		PAGE
Spirit of Camphor (A)	130	Taste and Taste Buds (S)	301
Spring Greens (S)	372	Tea and Coffee Habitués (S) ...	418
Spirochaetocides	206	Tenderizing Meat (S)	177
Spur Feterita	160	Testosterone Propionate (A) ...	267
Stabilization (S)	268	Textbook of Materia Medica, Pharmacology and Therapeutics (B)	484
Stabilization of Hydrogen Perox- ide for Pharmaceutical Pur- poses (A)	338	Thermolabile Substances (A) ...	123
Standard Chemical and Technical Dictionary (B)	340	Thermophilic Organisms in Sugar (A)	465
Starch Paste (A)	128	Thymol	230, 278, 386
Stearyl Alcohol (A)	76	Tobacco Tars Cause Cancer (S). 373	
Sterols and Related Compounds. 211		Tobacco With Lower Nicotine Content (S)	304
Stethographone	98	Toffana Water (E)	84
Stilbestrol (A)	174	Toxicity of Parachlormetacresol and Phenyl Mercuric Nitrate (A)	336
Strawberry Juice (S)	268	Tragic Neglect of Scientific Methods	19
Strychnine (A)	260	Traumatic Acid (S)	482
Strychnine Incompatible With Iodides (A)	266	Treatise on Pharmacy (B)	178
Sugar (S)	41	Treatment of Malaria With Mapharsen and Tryparsamide, Warning Against (A)	335
Sugars "Rare" (S)	41	Trehalose (S)	41
Sulfanilamide264, 303, 418, 481		Typhoid Statistics (S)	270
Sulfanilamido Derivations of Heterocyclic amines (A)	407	Ultraviolet Light	165
Sulfapyridine79, 297, 374, 477		Ultraviolet Radiation (A)	408
2-Sulfanilyl-Aminopyridine (Sulfa- pyridine), Absorption and Ex- cretion of (A)	219	United States Leadership in Chemistry (S)	372
Sulfonated Castor Oil Mixtures. 314		U. S. P. XI (E)	310
Sulfonated Oils	314	U. S. P. XI Supplement, Second. 319	
Sulfur, Color Reaction for (A). 336		U. S. P. XII (E)	313
Suppository Base, Water Soluble (A)	73	Vaccination (S)	133
Synthetic Camphor as a Pre- servative	387	Vaccines	147
Synthetic Gum (S)	458	Vacuum Drying of Extracts (A) 35	
Tannic Acid Gauze (A)	413	Vegetable Juices (E)	275
Tansy (E)	275	Veratrum Album	84
Tar Acid Oil, Phenol Coefficient of	317		

	PAGE		PAGE
Veratrum Viride	86	Watercress (S)	372
Veratrum Viride, Evaluation by		Watermelon Diuresis	68
Daphnia	426	Water in Oil Emulsifying Agents	
Vinyl Ether (A)	172	(A)	368
Viscosities of Medicinal Mineral		Weed Eradication (S)	457
Oils	139	Word Study (S)	41
Vital Stain (S)	481	World's Fair (E)	2
Vitamin A	37, 114, 166	Xenon	161
Vitamin B ₁	116, 373	X-rays	121, 331
Vitamin B ₂ as a New Nit and		Yellow Fever Virus (A)	78
Lice Foe (S)	417	Yohimbine	466
Vitamin C (S)	372, 458	Yuletide Ornament Sprightening	
Vitamin E (S)	374	(S)	480
Vitamin Discovery (S)	303	Zein and Casein as New Soap	
Vitamin K (S)	372	Additives (A)	183
Vitamin Review	209	Zinc Content of Commercial Sam-	
Vitamin T (S)	303	ples of Insulin (A)	126
Vitamin X, V and Z (S)	303	Zinc, Quinaldinic Acid as a Re-	
Vitamins (S)	131	agent for (A)	74
Vitamins and Hormones (B)	179	Zollickoffer Pharmacy (E)	44
Warning Against Treatment of			
Induced Malaria With Maphar-			
sen and Tryparsamide (A) ...	335		



Doctor says, "Come right in."

WE PAY HIM, BUT HE WORKS FOR YOU

● Receptionists in busy offices everywhere, welcome the Lilly man. Even though the doctor must be selective in his interviews, he wants to see the Lilly man, who through a sense of responsibility engendered by the experience of a long line of forebears, knows how to conserve the physician's time as well as his own.

Every successful interview conducted by the Lilly man in your territory means more prescriptions for your store. Every additional prescription means more profit at the end of the year. The Lilly man works diligently in your behalf every working day in the year, yet his services cost you nothing. Maintain adequate stocks. No man ever loses on a Lilly Product.



John S. Rowe completed his thirty-fourth year as a Lilly representative last October 1. Except for a brief period in western Maryland, the entire thirty-four years have been spent in North Carolina. Loved and respected by all who know him, Mr. Rowe now serves the drug trade in and around Hickory.

ELI LILLY AND COMPANY • Indianapolis, Indiana, U. S. A.

IF YOU DON'T KNOW YOUR VITAMINS —KNOW THE MAKER!

IRRADOL-A

Composition: A palatable syrupy-like liquid containing natural vitamin A; vitamin B₁ (thiamin chloride); vitamin B₂ or G (riboflavin); vitamin D; iron and ammonium citrate; and malt extract (non-diastatic).

Uses: As a comprehensive source of vitamins A, B₁, B₂ (G), and D, plus iron and malt extracts.

How Supplied: In 11-ounce wide-mouth bottles and 1-pound and 2¾-pound cylindrical glass jars.

NATOLA Liquid and Capsules

Composition: The vitamin A and D activity of Natola (Halibut Liver Oil Fortified with Natural Vitamin D) is entirely of natural origin and is derived from halibut liver oil, tuna liver oil, and other fish liver oils of high vitamin content. Each gram contains 55,000 units of vitamin A and 5500 units of vitamin D per gram, or 9400 units of vitamin A and 940 units of vitamin D per ten drops or capsule.

Uses: For administration of vitamins A and D in prevention and treatment of deficiencies of these factors.

How Supplied: Natola Liquid, in bottles of 10 cc. and 50 cc. with dropper cap; Natola Capsules (No. 217), in boxes of 25, 50, 100 and 250.

ABDOL WITH VITAMIN C CAPSULES (No. 218)

Composition: Each capsule contains 10,000 U.S.P. units of vitamin A; 1000 U.S.P. units of vitamin D; 200 International units of vitamin B₁ (Thiamin Chloride); 100 Gammas (40 Sherman units) of vitamin B₂ (G) (Riboflavin), and 500 International units of vitamin C (ascorbic acid)

Uses: To enable the physician to prescribe in a single preparation five important vitamin factors of standardized high potency—vitamins A, B₁, B₂ (G), C, and D.

How Supplied: Soluble gelatin capsules, packages of 25, 50, 100, and 250.

PARKE, DAVIS & COMPANY • Detroit, Mich.
The World's Largest Makers of Pharmaceutical and Biological Products

Keep this Profit-Making Display

**Near the
CASH
REGISTER**

**At the
TOBACCO
COUNTER**



'SUCRETS'

**Antiseptic Throat Lozenges—
Tested for Germicidal
Activity**

Dissolved slowly on the tongue, 'Sucret' troches produce a marked soothing effect on the mucous membrane of the mouth and throat . . . promptly allay that "scratchy" feeling.

Hexylresorcinol 'Sucrets' are convenient to carry. They are packed 24 in a neat metal box, each 'Sucret' individually wrapped. They may be carried loose in the handbag or pocket, or kept at the bedside.



'SUCRETS' TROCHES are a "pick-up item" for every customer with throat irritation! Displayed prominently near the cash register and at the tobacco counter, the 'Sucrets' counter container will make many reminder sales for you.

'Sucrets' are profitable, too, because every box is a quarter sale. And they bring satisfaction to your customer!



"For the Conservation of Life"

SHARP & DOHME

Pharmaceuticals

Mulford Biologicals

PHILADELPHIA

H O L A D I N

An Extract of the Entire Pancreas Gland

Holadin has long proved a definite resource in pancreatic disorders, diseases associated therewith. Holadin is put up in powder in 1 oz. and $\frac{1}{2}$ oz. vials, no "directions"; as commonly prescribed, the pharmacist may readily attach the usual prescription label.


FAIRCHILD BROS. & FOSTER

Holadin is also put up in capsules of 3 and 5 grains each.

New York

NOW READY!

A complete reference manual containing all essential information on dosage, therapeutic use, methods of administration etc., of all the biological products.



BIOLOGICAL PRODUCTS

by Dr. Louis Gershenfeld, P.D., B. Sc., Ph. M.

Professor of Bacteriology and Hygiene, Philadelphia College of Pharmacy and Science; Member of U. S. P. and N. F. Advisory Revision Committees; Consultant, Author, etc.

This volume represents a sound, accurate and concise source of information on antitoxins, serums, antigens, vaccines, toxoids, bacteriophage, allergens and other biological products in common use. Contained in 290 pages together with 40 illustrations, is the essential information that the bacteriologist, pharmacist, physician and student desires to have on hand for reference, on dosage, therapeutic use, methods of administration and the latest developments in connection with chemotherapy.

\$4.00

ADD 15c
For Postage
and Handling

ROMAINE PIERSON PUBLISHERS

99 NASSAU STREET
New York, N. Y.

